


Priority Principle™: Diagnosis/Condition




	Method	Dose	Notes
	Chronic	50 Hz x 3-5 minutes	URP, or PHT (shown) 50 Hz PHT (popliteal artery)
	Swelling	1000-3000 Hz 1-2 minutes each area	Ohshiro's Proximal Priority
	Inflammation	50 Hz x 2 minutes or TARGET/DOSE	
	Spasm	1000 Hz x 2 minutes per spasm	Pontinen's Principle
	Pain	500-1000 Hz for 3 minutes -----or 1000 Hz x 2-5 minutes	Systemic target (NRT) -----or Local targets
	Tissue Repair	5-250 Hz for 2 minutes or TARGET/DOSE -----and 50 Hz x 5 minutes	Local targets -----and Systemic target (PHT) at Popliteal artery
	ROM	1000 Hz x 1 minute	@2-3 location around the joint line

Priority Principle™: Diagnosis/Condition




Method	Dose	Notes
	50 Hz x 3-5 minutes	URP, or PHT (shown) 50 Hz PHT (popliteal artery)

Priority Principle™: Diagnosis/Condition




Method	Dose	Notes
 Swelling	1000-3000 Hz 1-2 minutes each area	Ohshiro's Proximal Priority
 Inflammation	50 Hz x 2 minutes or TARGET/DOSE	
 Spasm	1000 Hz x 2 minutes per spasm	Pontinen's Principle
 Pain	500-1000 Hz for 3 minutes -----or 1000 Hz x 2-5 minutes	Systemic target (NRT) -----or Local targets

Priority Principle™: Diagnosis/Condition



Method	Dose	Notes
	5-250 Hz for 2 minutes or TARGET/DOSE -----and 50 Hz x 5 minutes	Local targets -----and Systemic target (PHT) at Popliteal artery

Priority Principle™: Diagnosis/Condition



Method	Dose	Notes
	ROM	1000 Hz x 1 minute